فروویب®

آیرون پلی مالتوز ۵۰ میلی‌گرم/ ۵ میلی‌لیتر

۱۵۰ میلی لیتر شربت

با طعم کارامل

هر ۵ میلی‌لیتر حاوی:

آهـن (به فرم کمپلکس فریک هیدروکسـاید پلی مالتوز) ۵۰ میـلی گرم المنتال

مورد مصرف:

کمـک بـه تأمین آهن مورد نیاز بدن و جلوگیـری از کم‌خونی ناشی از فقر آهن.

این فرآورده صرفا مکمل اسـت و جهت پیشـگیری، تشـخیص یا درمان بیماری نمی باشـد.

مقدار و نحوه مصرف:

فروویب® را با توجه به نیاز فرد و مطابق دستور پزشک مصرف نمایید.

کودکان بالای ۲ سـال: ۲.۵ میلی‌لیتر روزانه (۲۵ میلی گرم آهن)

بزرگسـالان: ۵ میلی‌لیتر، یک تا دوبـار روزانه (۵۰ تا ۱۰۰ میلی گرم آهن)

قطره خوراکی فروویب® به منظور سهولت مصرف، جهت کمک به پیشگیری از کمبود آهن و کم‌خونی در کودکان زیر دو سال توصیه می گردد.

- جهـت تنظیـم دقیق مقدار مصـرف، از پیمانه مدرج داخل جعبه اسـتفاده نمایید.
- مقـدار مصـرف روزانه را می‌توان یک‌جا یا در چند نوبـت در طول روز مصرف نمود.
- جهـت کاهـش ناراحتی‌های گوارشی، می توان شـربت آهن را بلافاصله پـس از غذا مصرف نمود.
- به‌منظـور سـهولت مصرف، شـربت فروویب® را می‌توان با آب‌میوه یا غذای کـودک مخلوط نمود.
- قبـل از مصـرف، فرآورده را بـه خوبی تکان دهید تا محتویـات یکنواخت گردد.
- مقدار توصیه شـده را بدون مشـورت پزشک یا داروسـاز تغییر ندهید.

سایر ترکیبات:

سـوربیتول، سوکرالوز، سـاکارز، طعم‌دهنده کارامل، سدیم متیل پارابن، سدیم پروپیل پارابن، اتانول، آب.

عوارض جانبی:

مصرف مکمل های آهن ممکن اسـت باعث بروز عوارضی مانند درد شـکم، حالت تهوع، تغییر رنگ مدفوع، اسـهال، یبوسـت و بی اشتهایی گردد.

در صورت مشـاهده عوارض جانبی، با پزشک یا داروسـاز مشورت کنید.

موارد احتیاط و هشدارها:

- این فرآورده با دوز بالا می باشد و با توجه به اینکه مقدار آهن موجود در آن ٪ ۱۱/۱۱ بالاتر از حداکثر مقدار مصرف مجاز روزانه اسـت، می‌بایسـت حتما با مشـورت پزشـک یا داروساز مصرف شود.
- در صورت بروز حساسـیت، از مصرف فـرآورده خودداری نمایید.
- اگـر یـک نوبت را فرامـوش کردید، نوبت بعدی را طبق برنامـه مصرف نمایید.
- در دوران بارداری و شیردهی طبق دسـتور پزشـک یا داروسـاز مصرف نمایید.
- در صـورت وجـود مقادیر بالای آهـن در خون (ماننـد هموکروماتوزیس و همولیـز مزمن) از مصرف فـرآورده اجتناب نمایید.
- در صورت ابتلا به برخی اختلالات خونی مرتبط با آهن، مانند تالاسمی، آنمی سیدروبلاستیک یا آنمی ناشی از مسـمومیت با سـرب، قبل از مصرف فرآورده با پزشک یا داروسـاز مشورت نمایید.
- در صورت مصرف دارو یا مکمل های دیگر، قبل از مصرف این فرآورده با پزشک یا داروساز مشورت نمایید.

شرایط نگهداری:

- فرآورده از یخ‌زدگی محافظت شـود.
- فرآورده دور از دید و دسـترس کودکان نگهداری شـود.
- فرآورده در دمای کمتر از ۳۰ درجه سانتی‌گراد و داخل جعبه اصلی دور از نور مستقیم خورشید نگهداری شود.
- از مصرف فرآورده تاریخ گذشـته خودداری شـود.

بسته بندی:

فروویب® به صورت شـربت خوراکی در بطری شیشـه ای ۱۵۰ میلی‌لیتر، به همراه پیمانه مدرج و برگه راهنما عرضه می گردد.

دارنده پروانه سـاخت: برسیان دارو / ساخت ایران

مشـاوره و پاسخگوی علمی: ۰۲۱-۲۶۷۶۱۸۳۵

BARSIAN DAROU

Ferrovibe®

Iron Polymaltose 50 mg/5mL

150 mL Syrup

Caramel flavor

Each 5 mL contains:
Iron (as Ferric Hydroxide Polymaltose Complex) 50 mg elemental

Indication:
Ferrovibe® is used to support the body's iron requirements and prevent iron-deficiency anemia.
This product is a supplement and not intended to prevent, diagnose or treat any disease.

Dosage and Administration:
Take Ferrovibe® according to individual needs and as directed by a doctor.
Children over 2 years: 2.5 mL daily (25 mg iron)
Adults: 5 mL once or twice daily (50-100 mg iron)
Ferrovibe® oral drop is recommended to support the prevention of iron deficiency and anemia in children under two years of age, due to their ease of administration.

- To ensure accurate dosing, use the measuring cup provided in the package.
- The daily dose can be divided into separate doses or can be taken at once.
- To reduce gastrointestinal discomfort, the iron syrup may be taken immediately after a meal.
- Ferrovibe® can be mixed with fruit juices and with baby food.
- Shake well to get a homogeneous mixture before each use.
- Do not change the recommended dose without consulting a doctor or pharmacist.

Other Ingredients:
Sorbitol, Sucralose, Sucrose, Caramel flavor, Sodium methylparaben, Sodium propylparaben, Ethanol, Water.

Possible Side Effects:
Iron supplements may cause side effects, including abdominal pain, nausea, change in stool color, diarrhea, constipation and loss of appetite.
If you notice any side effects, consult a doctor or pharmacist.

Precautions and Warnings:

- This product is high dose and considering that the amount of iron in this product is 11.11% higher than upper intake level, it must be used as prescribed by a doctor or a pharmacist.
- Do not use this product if you are allergic to any of its ingredients.
- If you miss a dose, continue with your regular dosing schedule.
- Consult your doctor or pharmacist before use if you are pregnant or breastfeeding.
- Do not use this product if you have high iron levels in the blood, such as in hemochromatosis or chronic hemolysis.
- In case of iron-related hematologic disorders, such as thalassemia, sideroblastic anemia, or anemia associated with lead poisoning, consult a doctor or pharmacist before using this product.
- If you are taking other medications, you should consult with a doctor or pharmacist before using this product.

Storage Conditions:

- Do not freeze.
- Keep out of reach and sight of children.
- Store below 30°C in the original package to protect from direct sunlight.
- Do not use after the expiry date.

Packaging:
Each package of Ferrovibe® contains a 150 mL bottle of syrup with a measuring cup and a leaflet.

License Holder: Barsian Darou Co. / Made In Iran
Medical Advice and Support: +982126761835